

DIM Palmetto

Prostate Formula



Item # 74190
60 Softgels

The Possible DIM Palmetto Prostate Formula, a Food Supplement

- Supplies nutrients important for the healthy function of the prostate gland
- May support and regulate the effect of endogenous hormones on the prostate gland
- May promote the conversion of oestrogen to its beneficial, protective 2-hydroxyestrone metabolites and reduce production of genotoxic 16 α -hydroxyestrone
- Supports healthy detoxification enzyme systems

Description

DIM Palmetto Prostate Formula contains **BioResponse DIM™**, a unique formulation containing pure diindolylmethane, an indole. Indoles are plant compounds with health promoting properties, and are found in cruciferous vegetables such as broccoli, cabbage, cauliflower and Brussels sprouts. DIM® (diindolylmethane) has been shown to help regulate and promote a more efficient metabolism of oestrogen, and an optimal ratio of oestrogen metabolites important for male as well as female health. This formula combines DIM® with other nutrients to nutritionally support male reproductive health.

DIM Palmetto Prostate Formula includes a stable, bioavailable form of DIM®, made possible through a proprietary delivery system, that is co-solubilised with phosphatidylcholine, and microencapsulated in starch particles. Xenoestrogenic compounds, such as organochlorine pesticides, can significantly disrupt healthy oestrogen metabolism. These oestrogen disruptors alter oestradiol hydroxylation metabolism producing a higher ratio of the genotoxic 16 α -hydroxyestrone (16 α -OHE1) to the safer and weaker oestrogenic 2-hydroxyestrone (2-OHE1). DIM® can promote increased levels of the protective hydroxylated oestrogen 2-OHE1. The mechanisms for DIM®'s health benefits primarily involve the induction of mixed function oxidases and phase II detoxification enzyme systems by the binding and activation of the arylhydrocarbon receptor. Some have suggested that DIM® may also positively affect cellular signaling pathways.

Zinc is necessary for the functioning of over 300 enzymes, playing a crucial role in many biological processes. The epithelial cells of the

prostate gland accumulate higher zinc levels than any soft tissue in the male body, and zinc is necessary for the development of sperm. Research suggests that zinc may support prostate health through its participation in the regulation of 5-alpha-reductase. Zinc is also involved in the proper metabolism of unsaturated fatty acids.

Until 1936, pumpkin seeds were listed in the United States Pharmacopoeia. In recent decades, pumpkin seed oil has been utilised for prostate support, often used in conjunction with saw palmetto berry extract, and human and animal studies show **pumpkin seed oil** may also support the function of the bladder and urethra. Active ingredients include the essential fatty acids linoleic acid and oleic acid, tocopherols, amino acids, minerals, phytosterols, porphyrins and carotenoids including lutein.

We use a carbon dioxide supercritical solvent-free fluid extract of saw palmetto, which contains important essential fatty acids and phytosterols. Studies show that **saw palmetto** can reduce binding of dihydrotestosterone (DHT) in the prostate through inhibition of nuclear receptors. It also potentially inhibits the action of 5-alpha-reductase, the enzyme needed for the conversion of testosterone into DHT. Clinical trials have demonstrated that saw palmetto berry extract may support prostate health, while also showing no evidence that it interferes with measurement of serum prostate specific antigen (PSA), a marker used in monitoring the health of the prostate gland. There is also evidence that saw palmetto can play a role in healthy oestrogen metabolism.

Stinging Nettle root has been studied extensively regarding its detoxifying qualities and its support

of prostate function. It has potential to beneficially help regulate the effect of endogenous hormones, such as testosterone, dihydrotestosterone and oestrogen, on the prostate gland. Nettle root contains scopoletin, sterols, fatty acids, polysaccharides and lectins.

Beta-sitosterol is a sterol, a plant fat ("phytosterol") that closely resembles cholesterol. Phytosterols have been studied for decades and much is known about them. Beta-sitosterol has been widely researched and shown to support immunity, healthy cholesterol within normal levels and blood sugar within normal

levels. Three double-blind clinical trials of beta-sitosterol have demonstrated it has potential support for healthy prostate function.

Lycopene is a pigment that gives plants such as tomatoes, guava, watermelon and pink grapefruit their red hue. As an antioxidant carotenoid, it has been found to have protective effects for the vascular system and the eyes. Recent studies show that supplemental lycopene may support prostate function.

Serving size: 2 Softgels

Servings per container: 30

Amount per serving:

| | |
|--|--------|
| Calories | 10 |
| Calories from Fat | 10 |
| Total Fat | 1 g |
| Zinc (as Zinc Citrate) | 10 mg |
| Pumpkin Seed Oil | 893 mg |
| Saw Palmetto (Berries) Extract (Standardized to 85-95% Fatty Acids) | 320 mg |
| BioResponse DIM™ (A patented Diindolylmethane complex: starch, DIM (25% min.), Vitamin E succinate, phosphatidylcholine (soy), silica) | 300 mg |
| Stinging Nettle (Root) Extract 16:1 | 300 mg |
| Beta-Sitosterol | 120 mg |
| Lycopene | 15 mg |

Other ingredients: Gelatine, glycerine, water, yellow beeswax, carob extract, zinc oxide.

Suggested Use: As a food supplement, 2 softgels daily with food, or as directed by a healthcare practitioner.

Warning: Do not use this product if you are pregnant, lactating, or using birth control pills. Persons taking prescription medications should consult a healthcare professional before use. Harmless changes in urine colour may occur. Keep out of reach of children.

References

- | | |
|--|--|
| Berges RR, Kassen A, Senge T. BJU Int 2000;85:842-6. | Marks LS, Partin AW, Epstein JI, et al. J Urol 2000;163:1451-6. |
| Bonnesen C, Eggleston IM, et al. Cancer Res 2001;61:6120-30. | Shertzer HG, Senft AP. Drug Metabol Drug Interact 2000;17:159-88. |
| Chang YC, Riby J, Chang GH, et al. Biochem Pharmacol 1999;58:825-34. | Wilt T, Ishani A, MacDonald R, et al. Cochrane Database Syst Rev. 2000;(2):CD001043. Review. |
| Gerber GS. J Urol 2000;163:1408-12. | Krzeski T, et al. Clin Ther. 1993;15(6):1011-20. |
| Goepel M, Hecker U, Krege S, et al. Prostate 1999;38:208-15. | Wagner H, et al. Planta Med. 1989;55(5):452-54. |
| Koch E. Planta Med 2001;67:489-500. | Obertreis B, et al. Arzneim-Forsch/Drug Res. Jan1996;46(1):52-56. |
| Liang JY, Liu YY, Zou J, et al. Prostate 1999;40:200-7. | |
| Lu QY, Hung JC, Heber D, et al. Cancer Epidemiol Biomarkers Prev 2001;10:749-56. | |